

Markov Chain with discrete time and
discrete state space. $n \in \{0, 1, 2, \dots\}$

the value taken by MC
is at most countably many, state space S is
countable

$\{X_n\}$ is Markov chain, Markov property:

$$\begin{aligned} & \mathbb{P}(\underbrace{X_{n+1}=i_{n+1}}_{\text{future}} \mid \underbrace{X_n=i_n}_{\text{present}}, \underbrace{X_{n-1}=i_{n-1}, \dots, X_0=i_0}_{\text{past}}) \\ &= \mathbb{P}(\underbrace{X_{n+1}=i_{n+1}}_{\text{future}} \mid \underbrace{X_n=i_n}_{\text{present}}) \end{aligned}$$

$\Downarrow$

Transition probability matrix P .

$$P_{ij} = \mathbb{P}(X_{n+1}=j \mid X_n=i)$$

transition from i to j
(one-step)

$$(P^k)_{ij} = \mathbb{P}(X_{n+k}=j \mid X_n=i)$$

prob transition from i to j
(k -step)

e.g. $\{X_n\}$ MC, state space $S = \{1, 2, 3\}$.

$$P = \begin{pmatrix} 0.1 & 0.1 & 0.8 \\ 0.2 & 0.4 & 0.4 \\ 0.3 & 0.7 & 0 \end{pmatrix}$$

initial dist $\alpha^T = (0.1, 0.6, 0.3)$. Compute.

(a): $IP(X_7=1 | X_6=3) = P_{31} = \boxed{0.3}$

(b): $IP(\underbrace{X_9=3}_{\text{future}} | \underbrace{X_4=1, X_5=2}_{\text{past}}, \underbrace{X_7=1}_{\text{present}})$

Markov
property

$= IP(X_9=3 | X_7=1)$ (two-step transition prob from 1 to 3)

$$= (P^2)_{1,3} = \boxed{0.12}$$

$$P^2 = \begin{pmatrix} 0.27 & 0.61 & 0.12 \\ 0.22 & 0.46 & 0.32 \\ 0.17 & 0.31 & 0.52 \end{pmatrix}$$

$$(c): IP(X_0=2 | X_1=3)$$

$$\text{Apply Bayes thm} = \frac{IP(X_1=3 | X_0=2) \cdot IP(X_0=2)}{IP(X_1=3)}$$

$$= \frac{P_{23} \cdot 0.6}{IP(X_1=3)}$$

dist of X_1 is $\alpha^T \cdot P$

$$\begin{aligned} IP(X_1=j) &= \sum_i \underbrace{IP(X_0=i)} \cdot \underbrace{IP(X_1=j | X_0=i)} \\ &= \sum_i (\alpha^T)_i \cdot P_{ij} = (\alpha^T \cdot P)_j \end{aligned}$$

$$\alpha^T \cdot P = (0.1 \quad 0.6 \quad 0.3) \cdot \begin{pmatrix} 0.1 & 0.1 & 0.8 \\ 0.2 & 0.4 & 0.4 \\ 0.3 & 0.7 & 0 \end{pmatrix}$$

$$IP(X_1=3) = 0.1 \times 0.8 + 0.6 \times 0.4 + 0.3 \times 0$$

$$= 0.08 + 0.24 = 0.32$$

$$IP(X_0=2 | X_1=3) = \frac{0.4 \times 0.6}{0.32} = \frac{24}{32} = \boxed{\frac{3}{4}}$$

(d): $1EX_2$

transit twice
from X_0
to X_2 .

dist of X_2 is given by $\alpha^T \cdot P^2$

$$\alpha^T \cdot P^2 = (0.1 \quad 0.6 \quad 0.3) \begin{pmatrix} 0.27 & 0.61 & 0.12 \\ 0.22 & 0.46 & 0.32 \\ 0.17 & 0.31 & 0.52 \end{pmatrix}$$

$$= (0.21, 0.43, 0.36)$$

$$\begin{array}{ccc} \uparrow & \uparrow & \uparrow \\ 1P(X_2=1) & 1P(X_2=2) & 1P(X_2=3) \end{array}$$

$$1EX_2 = \sum_{i=1}^3 i \cdot 1P(X_2=i)$$

$$= 1 \times 0.21 + 2 \times 0.43 + 3 \times 0.36 = \boxed{2.15}$$

e.g: P always stochastic matrix (its row always sum up to 1)

P is called doubly stochastic if its columns also sum up to 1. (each row & each column sum up to 1)

Let $\{X_n\}$ be MC with state space $S = \{1, 2, \dots, k\}$, and its one-step transition matrix P is doubly stochastic, its initial dist is uniform on S .

Show that $\forall n, X_n$ is always uniform on S .

Pf: Apply induction,

①: For $n=0$, X_0 is uniform on S . ✓

②: Assume X_k is uniform on S , want to prove X_{k+1} is uniform on S .

$\forall i, IP(X_{k+1}=i)$ (law of total prob)

$$= \sum_{j=1}^k \underbrace{IP(X_k=j)}_{\frac{1}{k}} \cdot \underbrace{IP(X_{k+1}=i | X_k=j)}_{P_{ji}}$$


(induction assumption)

$$= \frac{1}{k} \underbrace{\sum_{j=1}^k P_{ji}}_{\substack{\text{sum of} \\ \text{the } i\text{-th column} \\ \text{of } P=1 \\ \text{(doubly stochastic)}}} = \frac{1}{k} \Rightarrow X_{k+1} \text{ is uniform on } \mathcal{S},$$

finishes the induction. ✓

Transition graph:

e.g:



what is $P_{ad} = \frac{2}{1+3+2} = \frac{2}{6}$

(normalize rates $\Rightarrow$ transition prob) $= \frac{1}{3}$

$$P_{bc} = \frac{4}{1+4+4+2} = \frac{4}{11}$$

$$P_{ij} = \frac{r_{ij}}{\sum_{k \in S} r_{ik}}$$