

Markov chain $\Rightarrow$ state space $\mathcal{S}$

each state $s \in \mathcal{S}$ $\begin{cases} \text{recurrent state} \\ \text{transient state} \end{cases}$
state classification

Recurrence:

Def: state s is called recurrent if

$$\mathbb{P}_{X_0=s}(T_s < \infty) = 1$$

start the MC
at state s

$T_s \triangleq \inf\{n \geq 1 : X_n = s\}$
is the first time that MC
hits state s (except time 0)

state s is recurrent iff starting from
state s , with probability 1, we can come back
to state s for infinitely many times.

Intuitively, Markov property means that the process is memoryless. so we can restart the process at any time.

If state s is recurrent, starting from $X_0 = s$, returning back to s cost finitely many time.


By Markov property, we act as if we restart the process after it hits s , so it only takes finitely many time until the second hit to s .

Do this for infinitely many times, as $\text{time} \rightarrow \infty$, each hit to state s cost finite time

$\Downarrow$

infinitely many visits to state s .

e.g: $\{X_n\}$ MC,



(a): Write transition matrix

state space $S = \{A, B, C, D, E, F, G\}$

	A	B	C	D	E	F	G
A		1					
B	$\frac{1}{2}$	$\frac{1}{2}$					
C			$\frac{1}{2}$	$\frac{1}{2}$			
D			1				
E			0.1			0.9	
F							1
G					1		

(b): Is there a limiting distribution?

$$\lim_{n \rightarrow \infty} \frac{\alpha^T P^n}{(x_n)}$$

Since not all states are communicating with each other, there's no single limiting dist, i.e. limiting dist depends on the initial dist α^T .

If we restrict to MC with state $\{A, B\}$ or the MC with state $\{C, D, E, F, G\}$, there is a single limiting distribution.

(c): Determine the set of stationary dist if exists

$$\pi^T \cdot P = \pi^T$$

$$(\pi_1, \dots, \pi_7) \cdot P = (\pi_1, \dots, \pi_7)$$

$$\left\{ \begin{array}{l} \frac{1}{2} \pi_2 = \pi_1 \\ \pi_1 + \frac{1}{2} \pi_2 = \pi_2 \quad (\times \text{ redundant}) \\ \frac{1}{2} \pi_3 + \pi_4 + \cancel{0.1 \pi_5} = \pi_3 \quad (\times \text{ redundant}) \\ \quad \quad \quad (\pi_5 = 0) \\ \frac{1}{2} \pi_3 = \pi_4 \\ \pi_7 = \pi_5 \\ 0.9 \pi_5 = \pi_6 \\ \pi_6 = \pi_7 \end{array} \right\} \quad \begin{array}{l} \pi_5 = \pi_7 = \pi_6 = 0.9 \pi_5 \\ \downarrow \\ \pi_5 = 0, \pi_6 = \pi_7 = 0. \end{array}$$

$$\left\{ \begin{array}{l} \pi_1 = \frac{1}{2} \pi_2 \\ \pi_4 = \frac{1}{2} \pi_3 \\ \pi_1 + \pi_2 + \pi_3 + \pi_4 = 1 \end{array} \right.$$

$$\pi_4 = a \text{ as free variable, } \pi_3 = 2a, \quad \left\{ \begin{array}{l} \pi_1 = \frac{1}{2} \pi_2 \\ \pi_1 + \pi_2 = 1 - 3a \end{array} \right.$$
$$\frac{3}{2} \pi_2 = 1 - 3a, \quad \pi_2 = \frac{2(1-3a)}{3}, \quad \pi_1 = \frac{1-3a}{3}$$

So: if π is stat dist,

$$\pi = \begin{bmatrix} \frac{1}{3}(1-3a) \\ \frac{2}{3}(1-3a) \\ 2a \\ a \\ 0 \\ 0 \\ 0 \end{bmatrix}$$

$$\begin{pmatrix} a \geq 0, \\ 1-3a \geq 0 \end{pmatrix}$$


$\Downarrow$

$$\left(0 \leq a \leq \frac{1}{3} \right)$$

here stat dist is not unique, since transition graph has two connected components.

these states are transient

If S transient, it must have value 0 in stationary dist.



(d): communication class

$\{A, B\}, \{C, D\}, \{E, F, G\}$

(e): recurrent / transient states

$\{A, B\}$ is closed $\Rightarrow$ A recurrent, B recurrent
 closed communication class
 with finitely many states is recurrent!


(all states in the same comm class
 has the same recurrence property)

$\{C, D\}$ is closed $\Rightarrow$ C, D recurrent.

$\{E, F, G\}$ is not closed $\Rightarrow$ E, F, G transient.

SRW $\Rightarrow$ MC

symmetric SRW $S = \mathbb{Z}$



since symmetric, there's no trend of going left/right for MC, recurrent.

asymmetric SRW p : prob of going right
($p \neq \frac{1}{2}$)


since asymmetric, there's a clear trend that it's either going left or going right, so starting from 0, we cannot expect to return to 0 again in finite time with prob 1
so it's transient.

$\{X_n\}$: MC , irreducible, closed , finitely many states



recurrent

$\{X_n\}$:



if we start the chain at 0,

$$X_0 = 0, X_1 = 1, X_2 = 0, X_3 = 1$$



first hitting time
back to 0,
so $T_0 = 2$.

$$\mathbb{P}_{X_0=0}(T_0 < \infty) = 1$$

by def, $\{X_n\}$ is recurrent.