

Markovian Diffusion $\{D_t\}$, cts sample path,

$$\begin{cases} \forall \varepsilon > 0, IP(|D_{t+h} - D_t| > \varepsilon | D_t = x) = o(h) \\ IE(D_{t+h} - D_t | D_t = x) = \underbrace{a(t, x)}_{\text{drift}} h + o(h) \quad (h \rightarrow 0) \\ IE[(D_{t+h} - D_t)^2 | D_t = x] = \underbrace{b(t, x)}_{\text{diffusion}} h + o(h) \end{cases}$$

Generalization: $dD_t = a(t, D_t)dt + b(t, D_t)dW_t$
(SDE)

How does $\{D_t\}$ propagate w.r.t. time?

$$\underbrace{f(t, y | s, x)}_{\text{conditional pdf}} \triangleq \frac{\partial}{\partial y} IP(D_t \leq y | D_s = x)$$

conditional pdf

of D_t given $D_s = x$,

y is the space variable

$$\begin{cases} \text{forward eqn: } \frac{\partial f}{\partial t} = -\frac{\partial}{\partial y} [a(t, y) \cdot f] \\ \quad \text{(Fokker-Planck)} \\ \quad \quad \quad + \frac{1}{2} \frac{\partial^2}{\partial y^2} [b(t, y) \cdot f] \\ \text{backward eqn: } \frac{\partial f}{\partial s} = -a(s, x) \frac{\partial f}{\partial x} \\ \quad \text{(Backward Kolmogorov)} \\ \quad \quad \quad - \frac{1}{2} b(s, x) \frac{\partial^2 f}{\partial x^2} \end{cases}$$

SDE
$\Updownarrow$
PDE

e.g: BM, $a(t,x)=0$, $b(t,x)=1$, $W_0=0$

$$\text{forward eqn: } \begin{cases} \frac{\partial f}{\partial t} = \frac{1}{2} \frac{\partial^2 f}{\partial y^2} \\ f(0,y) = \delta_0(y) \end{cases} \quad (s=0, x=0, f=f(t,y))$$

heat equation,
fundamental solution as Gaussian kernel.

Check: $f(t,y) = \frac{1}{\sqrt{2\pi t}} e^{-\frac{y^2}{2t}}$ is the solution.

$$\frac{\partial f}{\partial t} = \frac{1}{\sqrt{2\pi}} \left(-\frac{1}{2} t^{-\frac{3}{2}} e^{-\frac{y^2}{2t}} + t^{-\frac{1}{2}} \cdot e^{-\frac{y^2}{2t}} \cdot \frac{y^2}{2t^2} \right)$$

$$\frac{\partial f}{\partial y} = \frac{1}{\sqrt{2\pi t}} e^{-\frac{y^2}{2t}} \cdot \left(-\frac{y}{t} \right) = \frac{1}{\sqrt{2\pi}} t^{-\frac{3}{2}} (-y) \cdot e^{-\frac{y^2}{2t}}$$

$$\frac{\partial^2 f}{\partial y^2} = \frac{1}{\sqrt{2\pi}} t^{-\frac{3}{2}} \cdot \left(-e^{-\frac{y^2}{2t}} + \frac{y^2}{t} e^{-\frac{y^2}{2t}} \right)$$

$$\text{so } \frac{\partial f}{\partial t} = \frac{1}{2} \frac{\partial^2 f}{\partial y^2}, \quad f(0,y) = \lim_{t \rightarrow 0} \frac{1}{\sqrt{2\pi t}} e^{-\frac{y^2}{2t}} = \delta_0(y)$$

the solution f is actually pdf of $N(0,t)$
which matches $W_t \sim N(0,t)$.

eg: BM with drift $a(t, x) = m$, $b(t, x) = \sigma^2$,
 $X_t = \sigma W_t + mt + d$, so forward eqn:

$$\frac{\partial f}{\partial t} = -m \frac{\partial f}{\partial y} + \frac{\sigma^2}{2} \frac{\partial^2 f}{\partial y^2}$$

If an absorbing barrier is put up at 0,
 then it can be reflected in the boundary
 condition!

$$\begin{cases} f(0, y) = \delta_d(y) \\ f(t, 0) = 0 \end{cases} \rightarrow \text{if absorbed by 0, it's permanently 0!}$$

Let's solve this PDE with BC: pdf of $N(x+mt, t)$

first check $g(t, y|x) = \frac{1}{\sqrt{2\pi t}} e^{-\frac{(y-x-mt)^2}{2t}}$ is the
 solution to PDE for $\forall x$, but it does not
 satisfy the BC.

$\Downarrow$

$$f(t, y) = g(t, y|d) - e^{-2md} g(t, y|-d)$$

is the solution.

This is called "method of images", which is also the technique used in pricing barrier option, similar to reflection principle.

$T \triangleq$ first absorption time, time t $\begin{cases} \text{absorbed} \\ \text{or} \\ \text{has pdf } f \end{cases}$, so

$$IP(T \leq t) = 1 - \int_0^{+\infty} f(t, y) dy$$

$$= 1 - \Phi\left(\frac{d+mt}{\sqrt{t}}\right) + e^{-2md} \Phi\left(\frac{-d+mt}{\sqrt{t}}\right)$$

provides cdf of T .


$$IP(T < \infty) = e^{-2md} \text{ if } m > 0$$

$\Downarrow$

$$f_T(t) = \frac{d}{\sqrt{2\pi} t^{3/2}} e^{-\frac{(d+mt)^2}{2t}} \quad (t \geq 0)$$

What if there's a reflection barrier?

$f(t, y)$ now denotes the law of the reflected process at time t .



Then $\int_0^{+\infty} f(t, y) dy = 1$, $\forall t$, $\frac{\partial}{\partial t}$ on both sides,

$$0 = \int_0^{+\infty} \frac{\partial f}{\partial t} dy = \int_0^{+\infty} \left(-m \frac{\partial f}{\partial y} + \frac{\sigma^2}{2} \frac{\partial^2 f}{\partial y^2} \right) dy$$

int by parts

$$m f(t, 0) - \frac{\sigma^2}{2} \frac{\partial f}{\partial y}(t, 0)$$

So: forward eqn

$$\left\{ \begin{array}{l} \frac{\partial f}{\partial t} = -m \frac{\partial f}{\partial y} + \frac{\sigma^2}{2} \frac{\partial^2 f}{\partial y^2} \end{array} \right.$$

$$f(0, y) = \delta_d(y)$$

$$m \cdot f(t, 0) - \frac{\sigma^2}{2} \frac{\partial f}{\partial y}(t, 0) = 0$$

how to satisfy this?

Ansatz:

$$f(t, y) = g(t, y|d) + \underbrace{A \cdot g(t, y|-d)}_{\text{reflection}} + \underbrace{\int_{-\infty}^{-d} B(x) \cdot g(t, y|x) dx}_{\substack{\text{ots line of} \\ \text{reflections over} \\ (-\infty, -d)}}$$

↓

plug into the equation and calculate to get

$$A = e^{-2md}, \quad B(x) = -2m e^{2mx}$$

$$\begin{cases} \text{as } t \rightarrow \infty, f(t, y) \rightarrow 0 & \text{if } m \geq 0 \\ \text{as } t \rightarrow \infty, f(t, y) \rightarrow 2|m| e^{-2|m|x} & \text{if } m < 0 \end{cases}$$

$$\text{since } \int_{-\infty}^{-d} 2m e^{2mx} g(t, y|x) dx \quad u = \frac{x - y - mt}{\sqrt{t}}$$

$$= \int_{-\infty}^{-\frac{d+y+mt}{\sqrt{t}}} 2m e^{2my} \frac{1}{\sqrt{2\pi}} e^{-\frac{u^2}{2}} du$$

$$\rightarrow \begin{cases} 2|m| e^{-2|m|y} & \text{if } m < 0 \text{ as } t \rightarrow \infty. \\ 0 & \text{if } m \geq 0 \end{cases}$$